

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Goat F(ab)2 anti-Rat IgG (H+L)-FITC, MinX Bo,Ho,Hu,Ms,Rb,Sh, Polyclonal DNA-SEC-183860

| | |
|--------------------------|---|
| Artikelname | Goat F(ab)2 anti-Rat IgG (H+L)-FITC, MinX Bo,Ho,Hu,Ms,Rb,Sh, Polyclonal |
| Artikelnummer | DNA-SEC-183860 |
| Hersteller Artikelnummer | SEC-183860 |
| Alternativnummer | DNA-SEC-183860 |
| Hersteller | dianova |
| Wirt | Goat |
| Kategorie | Antikörper |
| Applikation | WB |
| Spezies Reaktivität | Rat |
| Immunogen | Rat IgG whole molecule |
| Konjugation | FITC |
| Produktbeschreibung | F(ab)2 Anti-Rat IgG (H&L) Antibody generated in goat detects rat IgG. Representing approximately 75% of serum immunoglobulins, IgG is the most abundant antibody isotype found in the circulation. IgG molecules are synthesized and secreted by plasma B ... |
| Klonalität | Polyclonal |
| Konzentration | 0.5 mg/mL |
| Isotyp | Ig |

| | |
|------------------------|--|
| Puffer | 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2 |
| Reinheit | This product was prepared from monospecific antiserum by immunoaffinity chromatography using Rat IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities, pepsin digestion and chromatographic separation. Assa |
| Formel | 20 mM K3PO4,150 mM NaCl,pH 7,2,lyophilisate,0,01% NaN3 |
| Target-Kategorie | Rat |
| Antibody Type | Polyclonal Antibody |
| Application Verdünnung | FLISA 1:10,000 - 1:50,000, FC 1:500 - 1:2,500, IF Microscopy 1:1,000 - 1:5,000 |
| Anwendungsbeschreibung | F(ab)2 Anti-Rat IgG (H&L) Antibody has been tested by western blot and is designed for immunofluorescence microscopy, fluorescence based plate assays (FLISA) and fluorescent western blotting. This product is also suitable for multiplex analysis, includin |