

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### Donkey Fab anti-Goat IgG (H+L)-unconj., MinX none, Polyclonal DNA-SEC-183927

|                          |   |
|--------------------------|---|
| Artikelname              | Donkey Fab anti-Goat IgG (H+L)-unconj., MinX none, Polyclonal   |
| Artikelnummer            | DNA-SEC-183927  |
| Hersteller Artikelnummer | SEC-183927  |
| Alternativnummer         | DNA-SEC-183927  |
| Hersteller               | dianova   |
| Wirt                     | Donkey  |
| Kategorie                | Antikörper  |
| Applikation              | WB, IHC, ELISA  |
| Spezies Reaktivität      | Goat  |
| Immunogen                | Goat IgG whole molecule   |
| Konjugation              | Unconjugated  |
| Produktbeschreibung      | Fab Anti-Goat IgG Antibody generated in donkey detects goat IgG. This product possesses the F(ab) region possessing the epitope-recognition site, both heavy and light chains of the antibody molecule are present. Secondary Antibodies are available in ... |
| Klonalität               | Polyclonal  |
| Konzentration            | 1.0 mg/mL   |
| Isotyp                   | Ig  |
| Puffer                   | 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2  |

|                        |  |
|------------------------|--|
| Reinheit               | This product was prepared from monospecific antiserum by immunoaffinity chromatography using Goat IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities, papain digestion and chromatographic separation. Ass |
| Formel                 | 20 mM K3PO4,150 mM NaCl,pH 7,2,sterile filtered,0,01% NaN3   |
| Target-Kategorie       | Goat   |
| Antibody Type          | Polyclonal Antibody  |
| Application Verdünnung | WB: 1:2,000 - 1:10,000   |
| Anwendungsbeschreibung | Fab Anti-Goat IgG Antibody has been tested by SDS-PAGE and is suitable for highly specific immunological methods requiring extremely low background levels, absence of F(c) mediated binding, lot-to-lot consistency, high titer and specificity.              |