

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### Rabbit Fab anti-Guinea Pig IgG (H+L)-HRPO, MinX none, Polyclonal DNA-SEC-183937

|                          |   |
|--------------------------|---|
| Artikelname              | Rabbit Fab anti-Guinea Pig IgG (H+L)-HRPO, MinX none, Polyclonal  |
| Artikelnummer            | DNA-SEC-183937  |
| Hersteller Artikelnummer | SEC-183937  |
| Alternativnummer         | DNA-SEC-183937  |
| Hersteller               | dianova   |
| Wirt                     | Rabbit  |
| Kategorie                | Antikörper  |
| Applikation              | WB, IHC, ELISA  |
| Spezies Reaktivität      | Guinea pig  |
| Immunogen                | Guinea Pig IgG whole molecule   |
| Konjugation              | HRP   |
| Produktbeschreibung      | Fab Anti-Guinea Pig IgG Peroxidase Antibody generated in rabbit detects guinea pig IgG. This product possesses the F(ab) region possessing the epitope-recognition site, both heavy and light chains of the antibody molecule are present. Secondary Antib... |
| Klonalität               | Polyclonal  |
| Konzentration            | 1.0 mg/mL   |
| Isotyp                   | Ig  |
| Puffer                   | 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2  |

|                        |   |
|------------------------|---|
| Reinheit               | This product was prepared from monospecific antiserum by immunoaffinity chromatography using Guinea Pig IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities, papain digestion and chromatographic separation |
| Formel                 | 20 mM K <sub>3</sub> PO <sub>4</sub> , 150 mM NaCl, pH 7,2, lyophilisate, 0,01% Gentamicin  |
| Target-Kategorie       | Guinea Pig  |
| Antibody Type          | Polyclonal Antibody   |
| Application Verdünnung | WB: 1:1,000 - 1:3,000   |
| Anwendungsbeschreibung | Suitable for immunoblotting (western or dot blot), ELISA, immunoperoxidase electron microscopy and immunohistochemistry as well as other peroxidase-antibody based enzymatic assays requiring extremely low background levels, absence of F(c) mediated binding |