

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### Rabbit Fab anti-Mouse IgG (H+L)-FITC, MinX none, Polyclonal DNA-SEC-183961

|                          |   |
|--------------------------|---|
| Artikelname              | Rabbit Fab anti-Mouse IgG (H+L)-FITC, MinX none, Polyclonal   |
| Artikelnummer            | DNA-SEC-183961  |
| Hersteller Artikelnummer | SEC-183961  |
| Alternativnummer         | DNA-SEC-183961  |
| Hersteller               | dianova   |
| Wirt                     | Rabbit  |
| Kategorie                | Antikörper  |
| Applikation              | FC, IF, FLISA   |
| Spezies Reaktivität      | Mouse   |
| Immunogen                | Mouse IgG whole molecule  |
| Konjugation              | FITC  |
| Produktbeschreibung      | Fab Anti-Mouse IgG (H&L) Fluorescein Antibody generated in rabbit detects Mouse IgG. This product possesses the F(ab) region possessing the epitope-recognition site, both heavy and light chains of the antibody molecule are present. Secondary Antibodi... |
| Klonalität               | Polyclonal  |
| Konzentration            | 0.5 mg/mL   |
| Isotyp                   | Ig  |
| Puffer                   | 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2  |

|                        |  |
|------------------------|--|
| Reinheit               | This product was prepared from monospecific antiserum by immunoaffinity chromatography using Mouse IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities, papain digestion and chromatographic separation. As |
| Formel                 | 20 mM K3PO4,150 mM NaCl,pH 7,2,lyophilisate,0,01% NaN3   |
| Target-Kategorie       | Mouse  |
| Antibody Type          | Polyclonal Antibody  |
| Application Verdünnung | FLISA 1:10,000 - 1:50,000, FC 1:500 - 1:2,500, IF Microscopy 1:1,000 - 1:5,000   |
| Anwendungsbeschreibung | Suitable for immunomicroscopy and flow cytometry or FACS analysis as well as other antibody based fluorescent assays requiring extremely low background levels, absence of F(c) mediated binding, lot-to-lot consistency, high titer and specificity. This pro |