

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### Donkey Fab anti-Sheep IgG (H+L)-unconj., MinX none, Polyclonal DNA-SEC-183984

|                          |   |
|--------------------------|---|
| Artikelname              | Donkey Fab anti-Sheep IgG (H+L)-unconj., MinX none, Polyclonal  |
| Artikelnummer            | DNA-SEC-183984  |
| Hersteller Artikelnummer | SEC-183984  |
| Alternativnummer         | DNA-SEC-183984  |
| Hersteller               | dianova   |
| Wirt                     | Donkey  |
| Kategorie                | Antikörper  |
| Applikation              | WB, IHC, ELISA  |
| Spezies Reaktivität      | Sheep   |
| Immunogen                | Sheep IgG whole molecule  |
| Konjugation              | Unconjugated  |
| Produktbeschreibung      | Fab Anti-Sheep IgG (H&L) Antibody generated in donkey detects immunoglobulin g from sheep, both heavy and light chains of the antibody molecule are present. Each IgG has two antigen binding sites. Representing approximately 75% of serum immunoglobuli... |
| Klonalität               | Polyclonal  |
| Konzentration            | 0.9 mg/mL   |
| Isotyp                   | Ig  |
| Puffer                   | 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2  |

|                        |  |
|------------------------|--|
| Reinheit               | This product was prepared from monospecific antiserum by immunoaffinity chromatography using Sheep IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities, papain digestion and chromatographic separation. As |
| Formel                 | 20 mM K3PO4,150 mM NaCl,pH 7,2,sterile filtered,0,01% NaN3   |
| Target-Kategorie       | Sheep  |
| Antibody Type          | Polyclonal Antibody  |
| Application Verdünnung | WB: 1:2,000 - 1:10,000   |
| Anwendungsbeschreibung | Suitable for highly specific immunological methods requiring extremely low background levels, absence of F(c) mediated binding, lot-to-lot consistency, high titer and specificity.  |