

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Sheep anti Human secretory component (free and bound), conjugated with Biotin, Clone: [Polyclonal], Monoclonal NMB-SHAHU/SC/BIO

| | |
|----------------------------|---|
| Article Name | Sheep anti Human secretory component (free and bound), conjugated with Biotin, Clone: [Polyclonal], Monoclonal |
| Biozol Catalog Number | NMB-SHAHU/SC/BIO |
| Supplier Catalog Number | ShAHu/SC/Bio |
| Alternative Catalog Number | NMB-SHAHU/SC/BIO |
| Manufacturer | NordicMubio |
| Host | Sheep |
| Category | Antikörper |
| Application | DOT, ELISA, ICC, IHC-P, WB |
| Species Reactivity | Human |
| Conjugation | Biotin |
| Product Description | Tested in immunoelectrophoresis, double radial immunodiffusion and ELISA against a panel of appropriate secretions and purified Ig isotypes. The antiserum reacts with both bound secretory component (secretory IgA) and with the free SC present in huma... |
| Clonality | Monoclonal |
| Clone Designation | [Polyclonal] |
| Buffer | Biotin-coupled purified hyperimmune sheep IgG lyophilized from a solution in phosphate buffered saline (PBS, pH 7.2). No preservative added, as it may interfere with the antibody activity. It is reconstituted by adding 1 ml sterile distilled water, spun |

| | |
|-------------------|--|
| Source | Secretory component is present in human secretions bound to secretory IgA (sIgA) and in free form. Secretory IgA (sIgA) functions as a dimer or polymer and accounts for almost all specific mucosal antibody activity. A molecule of sIgA is made up of two m |
| Application Notes | ELISA, Immunocytochemistry, Immunohistochemistry (paraffin), Dot blot, Immunoblotting. |