

Diagnostica Vertrieb GmbH, Leipziger Straße 4

85386 Eching, Germany

Telephone: +49 (0)89 3799666-6 | **Fax:** +49 (0)89 3799666-99

E-Mail: info@biozol.de

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Anti-Heme oxygenase 2/HMOX2 Antibody . Tested in IHC, WB applications. This antibody reacts with Human., Unconjugated, Rabbit, Polyclonal NBS-ABB-YECPDZ-100

| Article Name | Anti-Heme oxygenase 2/HMOX2 Antibody . Tested in IHC, WB applications. This antibody reacts with Human., Unconjugated, Rabbit, Polyclonal |
|----------------------------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Biozol Catalog Number | NBS-ABB-YECPDZ-100 |
| Supplier Catalog Number | ABB-YECPDZ-100 |
| Alternative Catalog Number | NBS-ABB-YECPDZ-100 |
| Manufacturer | Nordic BioSite |
| Host | Rabbit |
| Category | Antikörper |
| Application | IHC, WB |
| Species Reactivity | Human |
| Immunogen | E.coli-derived human HMOX2 recombinant protein (Position: S2-M316). Human HMOX2 shares 89% and 90% amino acid (aa) sequences identity with mouse and rat HMOX2, respectively. |
| Conjugation | Unconjugated |
| Product Description | Anti-Heme oxygenase 2/HMOX2 Antibody . Tested in IHC, WB applications. This antibody reacts with Human |
| Clonality | Polyclonal |
| Concentration | 500 ug/ml |
| Molecular Weight | 36033 MW |

| Isotype | IgG |
|--------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| NCBI | 3163 |
| UniProt | P30519 |
| Buffer | Each vial contains 5mg BSA, 0.9mg NaCl, 0.2mg Na2HPO4, 0.05mg NaN3. |
| Source | rabbit |
| Purity | Affinity Purified |
| Form | Lyophilized |
| Target | HMOX1 |
| Antibody Type | Primary Antibody |
| Application Dilute | IHC-P 0.5-1ug/ml, WB 0.1-0.5ug/ml |
| Application Notes | WB: The detection limit for HMOX2 is approximately 0.25ng/lane under reducing conditions. Tested Species: In-house tested species with positive results. By Heat: Boiling the paraffin sections in 10mM citrate buffer, pH6.0, for 20mins is required for the |