

Diagnostica Vertrieb GmbH, Leipziger Straße 4

85386 Eching, Germany

**Telephone:** +49 (0)89 3799666-6 | **Fax:** +49 (0)89 3799666-99

E-Mail: info@biozol.de

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

## **Product Datasheet**

## Influenza A H1N1 Hemagglutinin antibody, Clone: [AT1G7], Unconjugated, Mouse, Monoclonal NBS-AAT-UXM47T-100

| Article Name               | Influenza A H1N1 Hemagglutinin antibody, Clone: [AT1G7],<br>Unconjugated, Mouse, Monoclonal   |
|----------------------------|---|
| Biozol Catalog Number      | NBS-AAT-UXM47T-100  |
| Supplier Catalog Number    | AAT-UXM47T-100  |
| Alternative Catalog Number | NBS-AAT-UXM47T-100  |
| Manufacturer               | Nordic BioSite  |
| Host                       | Mouse   |
| Category                   | Antikörper  |
| Application                | ELISA, WB   |
| Species Reactivity         | Virus   |
| Immunogen                  | Recombinant Influenza A-H1N1/HA1 (18-344aa) purified from Baculovirus   |
| Conjugation                | Unconjugated  |
| Product Description        | Influenza virus, an enveloped virus of the Orthomyxoviridae family, has a unique capacity for genetic variation that is based in two molecular features of the virus family. First of all, the surface proteins of the virus are highly variable, able to |
| Clonality                  | Monoclonal  |
| Concentration              | 1 mg/ml   |
| Clone Designation          | [AT1G7]   |

| Isotype           | IgG1 k   |
|-------------------|--|
| NCBI              | 04238  |
| UniProt           | C7RYS4   |
| Buffer            | Liquid. In Phosphate-Buffered Saline (pH 7.4) with 0.02% Sodium Azide, 10% Glycerol.   |
| Source            | mouse  |
| Purity            | Protein G Purified   |
| Antibody Type     | Primary Antibody   |
| Application Notes | The antibody has been tested by ELISA, Western blot analysis to assure specificity and reactivity. Since application varies, however, each investigation should be titrated by the reagent to obtain optimal results. Recommended dilution range for Western b |