

Diagnostica Vertrieb GmbH, Leipziger Straße 4

85386 Eching, Germany

Telephone: +49 (0)89 3799666-6 | **Fax:** +49 (0)89 3799666-99

E-Mail: info@biozol.de

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Mouse anti Human CD14, conjugated with FITC, IgG2b, Clone: [FWKW], Monoclonal NMB-0142

| Article Name | Mouse anti Human CD14, conjugated with FITC, IgG2b, Clone: [FWKW], Monoclonal |
|----------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Biozol Catalog Number | NMB-0142 |
| Supplier Catalog Number | 0142 |
| Alternative Catalog Number | NMB-0142 |
| Manufacturer | NordicMubio |
| Host | Mouse |
| Category | Antikörper |
| Application | FC |
| Species Reactivity | Human |
| Conjugation | FITC |
| Product Description | Identification of human Monocyte/Macrophage cells expressing the 55 kDa M.W. surface antigen. CD14 is present on 70-93% of Monocyte/Macrophage cells in normal peripheral blood |
| Clonality | Monoclonal |
| Concentration | Titered for flow cytometry |
| Clone Designation | [FWKW] |
| Isotype | lgG2b |
| UniProt | P08571 |

| Buffer | Provided as solution in phosphate buffered saline with 0.08% sodium azide and 0.2% carrier protein |
|-------------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Purity | Protein A/G Chromatography |
| Formula | Provided as solution in phosphate buffered saline with 0.08% sodium azide and 0.2% carrier protein |
| Application Notes | PBMC: Add10 μ l of MAB/10 6 PBMC in 100 μ l PBS. Mix gently and incubate for 15 minutes at 2 to 8C. Wash twice with PBS and analyze or fix with 0.5% v/v of paraformaldehyde in PBS and analyze. WHOLE BLOOD: Add 10 μ l of MAB/100 μ l of whole blood. Mix gentl |