

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Mouse anti double-stranded RNA (J2, J5 and K1) Comparison Set, Clone: [J2 J5 and K1], Monoclonal NMB-10050100

Article Name	Mouse anti double-stranded RNA (J2, J5 and K1) Comparison Set, Clone: [J2 J5 and K1], Monoclonal
Biozol Catalog Number	NMB-10050100
Supplier Catalog Number	10050100
Alternative Catalog Number	NMB-10050100
Manufacturer	NordicMubio
Host	Mouse
Category	Antikörper
Application	DOT, ELISA, IAC, ICC, IHC
Product Description	Over the past decade our double-stranded RNA (dsRNA)antibodies have been used extensively to detect and characterise plant and animal viruses with dsRNA genomes or intermediates. In addition, the anti-dsRNA antibodies can be used as a diagnostic tool...
Clonality	Monoclonal
Concentration	Concentration after reconstitution: 1.00 mg/ml as determined by A280 nm (A280 nm = 1.47 corresponds to 1 mg/ml antibody).
Clone Designation	[J2 J5 and K1]
Isotype	IgG2a, IgG2b, IgG2a

Buffer	Mouse monoclonal antibody J2 recognises double-stranded RNA (dsRNA) provided that the length of the helix is greater than or equal to 40 bp. dsRNA-recognition is independent of the sequence and nucleotide composition of the antigen. All naturally occurri
Source	Female DBA/2 mice were injected intraperitoneally with a mixture of 50 ug L-dsRNA and 75 ug methylated bovine serum albumin, emulsified in complete Freund's adjuvant. After several boosts spleen cells were fused with Sp2/0-Ag14 myeloma cells to generate t
Purity	Gel electrophoretically pure IgG antibody.
Formula	The lyophilised samples should each be reconstituted with 100 µl sterile distilled water. The mAb will then be in PBS without any stabilisers or preservatives at a concentration of 1 mgr/ml. As a result of the lyophilisation procedure, the reconstituted
Application Notes	Mouse monoclonal antibodies J2, J5 and K1 can be used for ELISA, dsRNA-immunoblotting, immunoaffinity chromatography and in certain systems also for immunohistochemistry (see references). The optimum working dilution of each antibody for any specific app