

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

## Product Datasheet

### **CD15/ Lewis X antigen/ alpha-(1,3)-fucosyltransferase 4, IgM, Clone: [BRA4F1], Mouse, Monoclonal NMB-MUB0367P**

|                            |   |
|----------------------------|---|
| Article Name               | CD15/ Lewis X antigen/ alpha-(1,3)-fucosyltransferase 4, IgM, Clone: [BRA4F1], Mouse, Monoclonal  |
| Biozol Catalog Number      | NMB-MUB0367P  |
| Supplier Catalog Number    | MUB0367P  |
| Alternative Catalog Number | NMB-MUB0367P  |
| Manufacturer               | NordicMubio   |
| Host                       | Mouse   |
| Category                   | Antikörper  |
| Application                | FC, ICC, IHC-Fr, IHC-P  |
| Species Reactivity         | Human   |
| Product Description        | CD15 is a carbohydrate that is referred to as the Lewis X antigen, also known as alpha-(1,3)-fucosyltransferase 4, which is expressed on a variety of glycolipids and glycoproteins found on the cell surface of neutrophils, monocytes, eosinophils and n... |
| Clonality                  | Monoclonal  |
| Clone Designation          | [BRA4F1]  |
| Isotype                    | IgM   |
| UniProt                    | <a href="#">P22083</a>  |

|                   |   |
|-------------------|---|
| Buffer            | Each vial contains 100 µl 1mg/ml purified monoclonal antibody in PBS containing 0.09% sodium azide. The product is delivered in lyophilised format.   |
| Source            | BRA4F1 is a mouse IgM monoclonal antibody derived from the fusion of SP2.0-Ag14 mouse myeloma cells with splenocytes from a BALB/c mouse immunized with K562 cells.   |
| Formula           | Each vial contains 100 µl 1mg/ml purified monoclonal antibody in PBS containing 0.09% sodium azide. The product is delivered in lyophilised format.   |
| Application Notes | The BRA4F1 antibody is suitable for the detection of CD15 by flow cytometry, immunocytochemistry and immunohistochemistry of frozen and paraffin embedded tissues. Optimal antibody dilutions for the different applications should be determined by titration. |