

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Goat anti Bovine IgG (Fc specific), Clone: [Polyclonal], Monoclonal NMB-GAB/IGG(FC)

| Article Name | Goat anti Bovine IgG (Fc specific), Clone: [Polyclonal], Monoclonal |
|----------------------------|---|
| Biozol Catalog Number | NMB-GAB/IGG(FC) |
| Supplier Catalog Number | GAB/lgG(Fc) |
| Alternative Catalog Number | NMB-GAB/IGG(FC) |
| Manufacturer | NordicMubio |
| Host | Goat |
| Category | Antikörper |
| Product Description | The reactivity of the antiserum is restricted to the Fc part of the IgG molecule. In immunoelectrophoresis and radial immunodiffusion, using various antiserum concentrations against normal bovine serum a single precipitin line is obtained which shows |
| Clonality | Monoclonal |
| Clone Designation | [Polyclonal] |
| Buffer | Delipidated, heat inactivated, lyophilized, stable whole antiserum No preservative added. Total protein and IgG concentrations in the antiserum are comparable to those of pooled normal goat serum. No foreign proteins added. Reconstitute the lyophilized a |
| Source | Highly purified normal IgG isolated from pooled bovine serum. Freunds complete adjuvant is used in the first step of the immunization procedure. |

BIOZOL

| Formula | Delipidated, heat inactivated, lyophilized, stable whole antiserum No preservative added. Total protein and IgG concentrations in the antiserum are comparable to those of pooled normal goat serum. No foreign proteins added. |
|-------------------|---|
| Application Notes | Precipitation assays. In immunoelectrophoresis use 2 μ l serum or equivalent against 120 μ l antiserum. In double radial immunodiffusion (Ouchterlony) use a rosette arrangement with 10 μ l antiserum in 3 mm diameter center well and 2 μ l serum samples (neat |