

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

## **Product Datasheet**

## Goat anti Chicken IgM (Fc specific), conjugated with Horseradish peroxidase, Clone: [Polyclonal], HRP, Monoclonal NMB-GACH/IGM(FC)/PO

| Article Name               | Goat anti Chicken IgM (Fc specific), conjugated with Horseradish peroxidase, Clone: [Polyclonal], HRP, Monoclonal   |
|----------------------------|---|
| Biozol Catalog Number      | NMB-GACH/IGM(FC)/PO   |
| Supplier Catalog Number    | GACh/IgM(Fc)/PO   |
| Alternative Catalog Number | NMB-GACH/IGM(FC)/PO   |
| Manufacturer               | NordicMubio   |
| Host                       | Goat  |
| Category                   | Antikörper  |
| Conjugation                | HRP   |
| Product Description        | The reactivity of the antiserum is directed to the Fc subunit of the<br>IgM molecule which expresses strict isotypic (class) specificity. It<br>does not react with any non-Ig protein in chicken serum, as tested by<br>immunoelectrophoresis and double radial im     |
| Clonality                  | Monoclonal  |
| Clone Designation          | [Polyclonal]  |
| Buffer                     | Horseradish peroxidase-coupled purified hyperimmune goat IgG<br>lyophilized from a solution in phosphate buffered saline (PBS, pH<br>7.2). No preservative added, as it may interfere with the antibody<br>activity. It is reconstituted by adding 1 ml sterile distill |

| Source            | Purified normal IgM isolated from pooled chicken serum. Freunds<br>complete adjuvant is used in the first step of the immunization<br>procedure.   |
|-------------------|--|
| Formula           | Horseradish peroxidase-coupled purified hyperimmune goat IgG<br>lyophilized from a solution in phosphate buffered saline (PBS, pH<br>7.2). No preservative added, as it may interfere with the antibody<br>activity. |
| Application Notes | ELISA,Immunocytochemistry,Immunohistochemistry (paraffin),Dot<br>blot,Immunoblotting.  |