

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Goat anti Dog C3c, Clone: [Polyclonal], Monoclonal NMB-GAD/C3C

| | |
|----------------------------|--|
| Article Name | Goat anti Dog C3c, Clone: [Polyclonal], Monoclonal |
| Biozol Catalog Number | NMB-GAD/C3C |
| Supplier Catalog Number | GAD/C3c |
| Alternative Catalog Number | NMB-GAD/C3C |
| Manufacturer | NordicMubio |
| Host | Goat |
| Category | Antikörper |
| Product Description | In immunoelectrophoresis against fresh dog serum, a single precipitin line is obtained in the beta-1 region representing native C3. Against serum containing partly activated C3, a precipitin line is obtained which extends from the beta-1 into the alp... |
| Clonality | Monoclonal |
| Clone Designation | [Polyclonal] |
| UniProt | F1PGM1 |
| Buffer | Delipidated, heat inactivated, lyophilized, stable whole serum. No preservative added. Total protein and IgG concentrations in the antiserum are comparable to those of normal pooled goat serum. No foreign proteins added. Reconstitute the lyophilized anti |
| Source | C3c is the major fragment resulting from the C3 cleavage by C3 convertase and factor i. It is composed of an intact beta chain bound to two fragments of the alpha chain. The protein is isolated and purified from pooled normal dog serum by precipitation t |

| | |
|-------------------|--|
| Formula | Delipidated, heat inactivated, lyophilized, stable whole serum. No preservative added. Total protein and IgG concentrations in the antiserum are comparable to those of normal pooled goat serum. No foreign proteins added. |
| Application Notes | Precipitation assays. In immunoelectrophoresis use 2 μ l dog plasma or equivalent against 120 μ l antiserum. In double radial immunodiffusion use a rosette arrangement with 10 μ l antiserum in 3 mm diameter center well and 2 μ l plasma samples (neat and seri |