

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Goat anti Dog fibrinogen, Clone: [Polyclonal], Monoclonal NMB-GAD/FBG/7S

| | |
|----------------------------|--|
| Article Name | Goat anti Dog fibrinogen, Clone: [Polyclonal], Monoclonal |
| Biozol Catalog Number | NMB-GAD/FBG/7S |
| Supplier Catalog Number | GAD/Fbg/7S |
| Alternative Catalog Number | NMB-GAD/FBG/7S |
| Manufacturer | NordicMubio |
| Host | Goat |
| Category | Antikörper |
| Product Description | The reactivity of the antiserum is restricted to fibrinogen. In immunoelectrophoresis and radial immunodiffusion (Ouchterlony), using various antiserum concentrations against normal dog plasma a single precipitin line is obtained which shows a reacti... |
| Clonality | Monoclonal |
| Clone Designation | [Polyclonal] |
| UniProt | P68213 |
| Buffer | Purified hyperimmune IgG lyophilized from a solution in phosphate buffered saline (PBS, pH 7.2) No preservative added, as it may interfere with the antibody activity. Reconstitute the lyophilized antiserum by adding 1 ml sterile distilled water. Dilution |
| Source | Fibrinogen (clotting factor I) is a heat labile beta glycoprotein present in plasma. It is the precursor of fibrin, which is the key protein constituting the network of the blood clot. Thrombin converts fibrinogen to fibrin by limited proteolysis. Fibrin |

| | |
|-------------------|--|
| Formula | Purified hyperimmune IgG lyophilized from a solution in phosphate buffered saline (PBS, pH 7.2) No preservative added, as it may interfere with the antibody activity. |
| Application Notes | Indirect immunofluorescence,ELISA,Dot blot,Immunoblotting. |