

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Goat anti Dog IgA (Fc specific), conjugated with Biotin, Clone: [Polyclonal], Monoclonal NMB-GAD/IGA(FC)/BIO

Article Name	Goat anti Dog IgA (Fc specific), conjugated with Biotin, Clone: [Polyclonal], Monoclonal
Biozol Catalog Number	NMB-GAD/IGA(FC)/BIO
Supplier Catalog Number	GAD/IgA(Fc)/Bio
Alternative Catalog Number	NMB-GAD/IGA(FC)/BIO
Manufacturer	NordicMubio
Host	Goat
Category	Antikörper
Conjugation	Biotin
Product Description	The reactivity of the antiserum is directed to the Fc subunit of the IgA molecule which expresses strict isotypic (class) specificity. It does not react with any non-Ig protein in dog serum, as tested by immunoelectrophoresis and double radial immuno...
Clonality	Monoclonal
Clone Designation	[Polyclonal]
Buffer	Biotin-coupled purified hyperimmune goat IgG lyophilized from a solution in phosphate buffered saline (PBS, pH 7.2). No preservative added, as it may interfere with the antibody activity. IgG protein concentration 10 mg/ml. Biotin/IgG protein molar ratio

Source	Purified normal IgA isolated from pooled dog serum. Freund's complete adjuvant is used in the first step of the immunization procedure.
Formula	Biotin-coupled purified hyperimmune goat IgG lyophilized from a solution in phosphate buffered saline (PBS, pH 7.2). No preservative added, as it may interfere with the antibody activity. IgG protein concentration 10 mg/ml. Biotin/IgG protein molar ratio
Application Notes	ELISA, Immunocytochemistry, Immunohistochemistry (paraffin), Dot blot, Immunoblotting.