

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Goat anti Horse IgG1(T), Clone: [Polyclonal], Monoclonal NMB-GAHO/IGG1(T)

| | |
|----------------------------|--|
| Article Name | Goat anti Horse IgG1(T), Clone: [Polyclonal], Monoclonal |
| Biozol Catalog Number | NMB-GAHO/IGG1(T) |
| Supplier Catalog Number | GAHo/IgG1(T) |
| Alternative Catalog Number | NMB-GAHO/IGG1(T) |
| Manufacturer | NordicMubio |
| Host | Goat |
| Category | Antikörper |
| Product Description | The reactivity of the antiserum is directed to the Fc subunit of subclass IgG1. It does not react with IgG/Fab fragments, IgM and IgA or any non-Ig protein in horse serum, as tested by immunoelectrophoresis and double radial immunodiffusion. In preci... |
| Clonality | Monoclonal |
| Clone Designation | [Polyclonal] |
| Buffer | Delipidated, heat inactivated, lyophilized whole antiserum. No preservative added, as it may interfere with the antibody activity. Total protein and IgG concentration in the antiserum are comparable to those of pooled normal goat serum. No foreign protei |
| Source | Pools of purified normal IgG1 isolated from pooled serum from tetanus immunized horses. Freund's complete adjuvant is used in the first step of the immunization procedure. |

| | |
|-------------------|--|
| Formula | Delipidated, heat inactivated, lyophilized whole antiserum. No preservative added, as it may interfere with the antibody activity. Total protein and IgG concentration in the antiserum are comparable to those of pooled normal goat serum. No foreign protei |
| Application Notes | Precipitation assays. In immunoelectrophoresis use 2 µl or equivalent against 120 µl antiserum. In double radial immunodiffusion (Ouchterlony) use a rosette arrangement with 10 µl antiserum in a 3 mm diameter centre well and 2 µl serum samples (neat and |