

Diagnostica Vertrieb GmbH, Leipziger Straße 4

85386 Eching, Germany

Telephone: +49 (0)89 3799666-6 | **Fax:** +49 (0)89 3799666-99

E-Mail: info@biozol.de

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Goat anti Human ceruloplasmin, Clone: [Polyclonal], Monoclonal NMB-GAHU/CPL

| Article Name | Goat anti Human ceruloplasmin, Clone: [Polyclonal], Monoclonal |
|----------------------------|--|
| Biozol Catalog Number | NMB-GAHU/CPL |
| Supplier Catalog Number | GAHu/CpI |
| Alternative Catalog Number | NMB-GAHU/CPL |
| Manufacturer | NordicMubio |
| Host | Goat |
| Category | Antikörper |
| Product Description | The reactivity of the antiserum is restricted to alpha-2 ceruloplasmin. In immunoelectrophoresis and radial immunodiffusion (Ouchterlony), using various antiserum concentrations against fresh normal human plasma a single precipitin line is obtained w |
| Clonality | Monoclonal |
| Clone Designation | [Polyclonal] |
| UniProt | P00450 |
| Buffer | Delipidated, heat inactivated, lyophilized, stable whole antiserum No preservative added. Total protein and IgG concentrations in the antiserum are comparable to those of pooled normal goat serum. No foreign proteins added. Reconstitute the lyophilized a |
| Source | Alpha-2 ceruloplasmin is copper containing alpha-2 glycoprotein with a molecular weight of 160,000. Its mean concentration in plasma of adults is 32 mg/100ml. Elevated levels are observed in pregnant women, in patients with infectious diseases and the ne |

| Formula | Delipidated, heat inactivated, lyophilized, stable whole antiserum No preservative added. Total protein and IgG concentrations in the antiserum are comparable to those of pooled normal goat serum. No foreign proteins added. |
|-------------------|--|
| Application Notes | Precipitation assays. In immunoelectrophoresis use 2 μ l or equivalent against 120 μ l antiserum. In double radial immunodiffusion (Ouchterlony) use a rosette arrangement with 10 μ l in a 3 mm diameter center well and 2 μ l serum samples (neat and serially d |