

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Goat anti human Factor VIII related antigen (VWD), Clone: [Polyclonal], Monoclonal NMB-GAHU/FVIII VWD

Article Name	Goat anti human Factor VIII related antigen (VWD), Clone: [Polyclonal], Monoclonal
Biozol Catalog Number	NMB-GAHU/FVIII VWD
Supplier Catalog Number	GAHu/FVIII VWD
Alternative Catalog Number	NMB-GAHU/FVIII VWD
Manufacturer	NordicMubio
Host	Goat
Category	Antikörper
Product Description	The defined antibody reactivity is restricted to VWF. In immunoelectrophoresis, bi-dimensional electrophoresis and radial immunodiffusion (Ouchterlony) against normal plasma, a single precipitin line is obtained which shows a reaction of identity wit...
Clonality	Monoclonal
Clone Designation	[Polyclonal]
Buffer	Delipidated, heat inactivated, lyophilized, stable whole serum. No preservative added. IgG protein concentration in the antiserum is 10 mg/ml. No foreign proteins added. Reconstitute the lyophilized antiserum by adding 1 ml sterile distilled water.
Source	Human FVIII procoagulant activity (FVIIIIC) is carried by a polypeptide non-covalently bound to a large carrier molecule (unit molecular weight 250 kD) known as the Von Willebrand factor (VWF). FVIII/VWF exists in plasma as a series of polymers with molec

Formula	Delipidated, heat inactivated, lyophilized, stable whole serum. No preservative added. IgG protein concentration in the antiserum is 10 mg/ml. No foreign proteins added.
Application Notes	The antiserum concentration required in the gel is normally between 1 and 2%. In immunologic determinations of FVIII/VWF, plasma samples and all assay components must contain EDTA to stabilize the proteins The antiserum is standardized for use in the ele