

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Goat anti Human IgG (Fc specific), conjugated with FITC, Clone: [Polyclonal], Monoclonal NMB-SR/GAHU/IGG(FC)/FITC

| | |
|----------------------------|--|
| Article Name | Goat anti Human IgG (Fc specific), conjugated with FITC, Clone: [Polyclonal], Monoclonal |
| Biozol Catalog Number | NMB-SR/GAHU/IGG(FC)/FITC |
| Supplier Catalog Number | SR/GAHu/IgG(Fc)/FITC |
| Alternative Catalog Number | NMB-SR/GAHU/IGG(FC)/FITC |
| Manufacturer | NordicMubio |
| Host | Goat |
| Category | Antikörper |
| Application | ELISA, ICC, IF, IHC-Fr |
| Species Reactivity | Human |
| Conjugation | FITC |
| Product Description | The reactivity of the antiserum is directed to the Fc subunit of the IgG molecule, which expresses strict (class) specificity. In immunoelectrophoresis and radial in immunodiffusion using various antiserum concentrations against human serum, a single... |
| Clonality | Monoclonal |
| Clone Designation | [Polyclonal] |
| Buffer | FITC-coupled purified hyperimmune goat IgG lyophilized from a solution in phosphate buffered saline (PBS, pH 7.2). No preservative added, as it may interfere with the antibody activity. It is reconstituted by adding 1 ml sterile distilled water, spun dow |

| | |
|-------------------|---|
| Source | Pools of purified IgG isolated from human serum. Freund's complete adjuvant is used in the first step of the immunization procedure. |
| Formula | FITC-coupled purified hyperimmune goat IgG lyophilized from a solution in phosphate buffered saline (PBS, pH 7.2). No preservative added, as it may interfere with the antibody activity. |
| Application Notes | ELISA, Immunocytochemistry, Immunohistochemistry (frozen), (In)direct immunofluorescence. |