

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Goat anti Human IgG IgA IgM (Fc specific), Clone: [Polyclonal], Monoclonal NMB-GAHU/IG(FC)

| | |
|----------------------------|--|
| Article Name | Goat anti Human IgG IgA IgM (Fc specific), Clone: [Polyclonal], Monoclonal |
| Biozol Catalog Number | NMB-GAHU/IG(FC) |
| Supplier Catalog Number | GAHu/Ig(Fc) |
| Alternative Catalog Number | NMB-GAHU/IG(FC) |
| Manufacturer | NordicMubio |
| Host | Goat |
| Category | Antikörper |
| Species Reactivity | Human |
| Product Description | The reactivity of the antiserum is directed to the Fc subunits of the major immunoglobulins. It does not react with any non-Ig protein in human serum, as tested by immunoelectrophoresis and double radial immunodiffusion. In precipitating techniques a... |
| Clonality | Monoclonal |
| Clone Designation | [Polyclonal] |
| Buffer | Delipidated, heat inactivated, lyophilized stable whole serum No preservative added, as it may interfere with the antibody activity. No foreign protein added. Total protein and IgG concentration in the antiserum are comparable to those of pooled goat ser |

| | |
|-------------------|---|
| Source | Purified normal IgG, IgA and IgM fractions isolated from pooled human serum. Freund's complete adjuvant is used in the first step of the immunization procedure. |
| Formula | Delipidated, heat inactivated, lyophilized stable whole serum. No preservative added, as it may interfere with the antibody activity. No foreign protein added. Total protein and IgG concentration in the antiserum are comparable to those of pooled goat serum. |
| Application Notes | Precipitation assays. In immunoelectrophoresis use 2 μ l or equivalent against 120 μ l antiserum. In double radial immunodiffusion use a rosette arrangement with 10 μ l antiserum in a 3 mm diameter centre well and 2 μ l serum samples (neat and diluted) in 2 |