

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Goat anti Human IgG IgA IgM (Fc specific), conjugated with Horseradish peroxidase, Clone: [Polyclonal], HRP, Monoclonal NMB-GAHU/IG(FC)/PO

| | |
|----------------------------|---|
| Article Name | Goat anti Human IgG IgA IgM (Fc specific), conjugated with Horseradish peroxidase, Clone: [Polyclonal], HRP, Monoclonal |
| Biozol Catalog Number | NMB-GAHU/IG(FC)/PO |
| Supplier Catalog Number | GAHu/Ig(Fc)/PO |
| Alternative Catalog Number | NMB-GAHU/IG(FC)/PO |
| Manufacturer | NordicMubio |
| Host | Goat |
| Category | Antikörper |
| Conjugation | HRP |
| Product Description | The reactivity of the antiserum is directed to the Fc subunits of the major isotypes of the human immunoglobulin system. No reaction is obtained with purified IgG/Fab or any non-Ig protein of human serum, as tested by immunoelectrophoresis and double... |
| Clonality | Monoclonal |
| Clone Designation | [Polyclonal] |
| Buffer | Peroxidase-coupled purified hyperimmune goat IgG lyophilized from a solution in phosphate buffered saline (PBS, pH 7.2). No preservative added, as it may interfere with the antibody activity. It is reconstituted by adding 1 ml sterile distilled water. |

| | |
|-------------------|---|
| Source | Purified polyclonal human IgG, and IgA and IgM containing factions isolated from human serum. Freund's complete adjuvant is used in the first step of the immunization procedure. |
| Formula | Peroxidase-coupled purified hyperimmune goat IgG lyophilized from a solution in phosphate buffered saline (PBS, pH 7.2). No preservative added, as it may interfere with the antibody activity. |
| Application Notes | ELISA, Immunocytochemistry, Immunohistochemistry (paraffin), Dot blot, Immunoblotting. |