

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

## Product Datasheet

### Goat anti Human J chain of dimeric IgA, Clone: [Polyclonal], Monoclonal NMB-GAHU/JSET

|                            |  |
|----------------------------|--|
| Article Name               | Goat anti Human J chain of dimeric IgA, Clone: [Polyclonal], Monoclonal  |
| Biozol Catalog Number      | NMB-GAHU/JSET  |
| Supplier Catalog Number    | GAHu/J set   |
| Alternative Catalog Number | NMB-GAHU/JSET  |
| Manufacturer               | NordicMubio  |
| Host                       | Goat   |
| Category                   | Antikörper   |
| Product Description        | The reactivity of the antiserum is restricted to J chain as tested in immunoelectrophoresis and radial immunodiffusion. The antiserum shows a single precipitation reaction with totally reduced and alkylated polyclonal and monoclonal polymeric IgA and...  |
| Clonality                  | Monoclonal   |
| Clone Designation          | [Polyclonal]   |
| Buffer                     | Delipidated, heat inactivated, lyophilized, stable whole antiserum No preservative added. Total protein and IgG concentrations in the antiserum are comparable to those of pooled normal goat serum. No foreign proteins added. Reconstitute the lyophilized a |
| Source                     | Human J chain is a polypeptide folded within the polymeric structure of the immunoglobulin. J chain isolated from human polymeric IgA and IgM are identical by criteria of composition, peptide maps and antigenicity. Human J chain has been established as d |

|                   |  |
|-------------------|--|
| Formula           | Delipidated, heat inactivated, lyophilized, stable whole antiserum No preservative added. Total protein and IgG concentrations in the antiserum are comparable to those of pooled normal goat serum. No foreign proteins added.                                |
| Application Notes | Precipitation assays. In immunoelectrophoresis use 2 µl serum or equivalent against 120 µl antiserum. In double radial immunodiffusion use a rosette arrangement with 10 µl antiserum in 3 mm diameter centre well and 2 µl serum samples (neat and serially d |