

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

## Product Datasheet

### Goat anti Human J chain of dimeric IgA, conjugated with FITC, Clone: [Polyclonal], Monoclonal NMB-GAHU/J/FITC

|                            |   |
|----------------------------|---|
| Article Name               | Goat anti Human J chain of dimeric IgA, conjugated with FITC, Clone: [Polyclonal], Monoclonal   |
| Biozol Catalog Number      | NMB-GAHU/J/FITC   |
| Supplier Catalog Number    | GAHu/J/FITC   |
| Alternative Catalog Number | NMB-GAHU/J/FITC   |
| Manufacturer               | NordicMubio   |
| Host                       | Goat  |
| Category                   | Antikörper  |
| Conjugation                | FITC  |
| Product Description        | In immunoelectrophoresis and radial immunodiffusion (Ouchterlony), this immunoconjugate shows a single precipitation reaction with totally reduced and alkylated polyclonal and monoclonal polymeric IgA, secretory IgA and IgM. A reaction of complete id... |
| Clonality                  | Monoclonal  |
| Clone Designation          | [Polyclonal]  |
| Buffer                     | FITC-coupled purified hyperimmune IgG lyophilized from a solution in phosphate buffered saline (PBS, pH 7.2) No preservative added, as it may interfere with the antibody activity.   |

|                   |  |
|-------------------|--|
| Source            | Human J chain is a polypeptide folded within the structure of the polymeric immunoglobulin, resulting in a very limited exposure of J chain antigenic determinants. The antiserum is raised against the isolated and purified J chain. J chains isolated from  |
| Formula           | FITC-coupled purified hyperimmune IgG lyophilized from a solution in phosphate buffered saline (PBS, pH 7.2) No preservative added, as it may interfere with the antibody activity.  |
| Application Notes | ELISA, Immunocytochemistry, Immunohistochemistry (frozen), (In)direct immunofluorescence. The lyophilized conjugate is shipped at ambient temperature and may be stored at +4C, prolonged storage at or below -20C. It is reconstituted by adding 1 ml sterile |