

Diagnostica Vertrieb GmbH, Leipziger Straße 4

85386 Eching, Germany

Telephone: +49 (0)89 3799666-6 | **Fax:** +49 (0)89 3799666-99

E-Mail: info@biozol.de

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Goat anti Human plasminogen, Clone: [Polyclonal], Monoclonal NMB-GAHU/PLG

Article Name	Goat anti Human plasminogen, Clone: [Polyclonal], Monoclonal
Biozol Catalog Number	NMB-GAHU/PLG
Supplier Catalog Number	GAHu/Plg
Alternative Catalog Number	NMB-GAHU/PLG
Manufacturer	NordicMubio
Host	Goat
Category	Antikörper
Product Description	The defined antibody reactivity is restricted to plasminogen as tested at the level of sensitivity of immuno-precipitation techniques. A single precipitin line is obtained in bidimensional electrophoresis, immunoelectrophoresis and double radial immu
Clonality	Monoclonal
Clone Designation	[Polyclonal]
UniProt	P00747
Buffer	Delipidated, heat inactivated, lyophilized, stable whole serum. No preservative added. Total protein and IgG concentrations in the antiserum are comparable to those of pooled normal goat serum. No foreign proteins added. Reconstitute the lyophilized anti
Source	Native plasminogen is a single polypeptide chain, synthesized in the liver. Its molecular weight has been reported to be 81,000 and 92,000. Part of the molecule contains the active serine esterase of plasmin. On activation it is converted to two polypept

Formula	Delipidated, heat inactivated, lyophilized, stable whole serum. No preservative added. Total protein and IgG concentrations in the antiserum are comparable to those of pooled normal goat serum. No foreign proteins added.
Application Notes	Precipitation assays. In electroimmunodiffusion the amount of antiserum required in the agarose gel is usually between 1 and 2% depending on the test arrangement.