

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

## **Product Datasheet**

## Goat anti Monkey C3c, Clone: [Polyclonal], Monoclonal NMB-GAMON/C3C/7S

| Article Name               | Goat anti Monkey C3c, Clone: [Polyclonal], Monoclonal   |
|----------------------------|---|
| Biozol Catalog Number      | NMB-GAMON/C3C/7S  |
| Supplier Catalog Number    | GAMon/C3c/7S  |
| Alternative Catalog Number | NMB-GAMON/C3C/7S  |
| Manufacturer               | NordicMubio   |
| Host                       | Goat  |
| Category                   | Antikörper  |
| Product Description        | In immunoelectrophoresis against fresh monkey serum, a single<br>precipitin line is obtained in the beta-1 region representing native<br>C3. Against serum containing partly activated C3, a precipitin line is<br>obtained which extends from the beta-1 into the      |
| Clonality                  | Monoclonal  |
| Clone Designation          | [Polyclonal]  |
| Buffer                     | Purified hyperimmune IgG lyophilized from a solution in phosphate<br>buffered saline (PBS, pH 7.2) No preservative added, as it may<br>interfere with the antibody activity. It is reconstituted by adding 1 ml<br>sterile distilled water, spun down to remove insolub |
| Source                     | C3c is the major fragment resulting from C3 cleavage by C3<br>convertase and factor I. It is composed of an intact beta chain bound<br>to two fragments of the alpha chain. C3c is isolated and purified from<br>pooled normal rhesus monkey serum. Freunds complete a  |

| Formula           | Purified hyperimmune IgG lyophilized from a solution in phosphate<br>buffered saline (PBS, pH 7.2) No preservative added, as it may<br>interfere with the antibody activity. |
|-------------------|--|
| Application Notes | Indirect immunofluorescence, ELISA, Dot blot, Immunoblotting.  |