

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Goat anti Monkey IgM (Fc specific), conjugated with Biotin, Clone: [Polyclonal], Monoclonal NMB-GAMON/IGM(FC)/BIO

| | |
|----------------------------|--|
| Article Name | Goat anti Monkey IgM (Fc specific), conjugated with Biotin, Clone: [Polyclonal], Monoclonal |
| Biozol Catalog Number | NMB-GAMON/IGM(FC)/BIO |
| Supplier Catalog Number | GAMon/IgM(Fc)/Bio |
| Alternative Catalog Number | NMB-GAMON/IGM(FC)/BIO |
| Manufacturer | NordicMubio |
| Host | Goat |
| Category | Antikörper |
| Conjugation | Biotin |
| Product Description | The reactivity of the antiserum is directed to the Fc subunit of the IgM molecule which expresses strict isotypic (class) specificity. It does not react with IgG, IgA and IgG/Fab or any non-Ig protein in monkey serum, as tested by immunoelectrophores... |
| Clonality | Monoclonal |
| Clone Designation | [Polyclonal] |
| Buffer | Biotin-coupled purified hyperimmune goat IgG lyophilized from a solution in phosphate buffered saline (PBS, pH 7.2). No preservative added, as it may interfere with the antibody activity. It is reconstituted by adding 1 ml sterile distilled water, spun d |
| Source | Purified IgM isolated from Rhesus monkey serum. Freund's complete adjuvant is used in the first step of the immunization procedure. |

| | |
|-------------------|---|
| Formula | Biotin-coupled purified hyperimmune goat IgG lyophilized from a solution in phosphate buffered saline (PBS, pH 7.2). No preservative added, as it may interfere with the antibody activity. |
| Application Notes | ELISA, Immunocytochemistry, Immunohistochemistry (paraffin), Dot blot, Immunoblotting. |