

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

## Product Datasheet

### Goat anti Monkey lactoferrin, Clone: [Polyclonal], Monoclonal NMB-GAMON/LFR

Article Name	Goat anti Monkey lactoferrin, Clone: [Polyclonal], Monoclonal
Biozol Catalog Number	NMB-GAMON/LFR
Supplier Catalog Number	GAMon/Lfr
Alternative Catalog Number	NMB-GAMON/LFR
Manufacturer	NordicMubio
Host	Goat
Category	Antikörper
Product Description	In immunoelectrophoresis against monkey milk a single precipitin line is obtained. The antiserum does not react with any other protein component of monkey serum or plasma. In precipitating techniques as immunoelectrophoresis, single and double radial...
Clonality	Monoclonal
Clone Designation	[Polyclonal]
Buffer	Delipidated, heat inactivated, lyophilized, stable whole serum. No preservative added. Total protein and IgG concentration in the antiserum are comparable to those of pooled normal goat serum. No foreign proteins added. Reconstitute the lyophilized artis
Source	Exocrine organs produce various secretions, each with its characteristic function. Proteins found in secretions may be divided into two groups: those specific for the particular secretion, and plasma proteins independent of the type of exocrine cells. La

Formula	Delipidated, heat inactivated, lyophilized, stable whole serum. No preservative added. Total protein and IgG concentration in the antiserum are comparable to those of pooled normal goat serum. No foreign proteins added.
Application Notes	Precipitation assays. In immunoelectrophoresis use 2 µl monkey serum against 120 µl antiserum. In double radial immunodiffusion (Ouchterlony) use a rosette arrangement with 10 µl antiserum in a 3 mm diameter center well and 10 µl of the antiserum in 3 mm