

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Goat anti Monkey secretory component (free and bound), Clone: [Polyclonal], Monoclonal NMB-GAMON/SC

Article Name	Goat anti Monkey secretory component (free and bound), Clone: [Polyclonal], Monoclonal
Biozol Catalog Number	NMB-GAMON/SC
Supplier Catalog Number	GAMon/SC
Alternative Catalog Number	NMB-GAMON/SC
Manufacturer	NordicMubio
Host	Goat
Category	Antikörper
Application	IP
Species Reactivity	Monkey
Product Description	Secretory IgA (sIgA) functions as a dimer or polymer and accounts for almost all specific mucosal antibody activity. A molecule of sIgA is made up of two molecules of IgA, one J chain and one SC (MW 65,000). The dimer IgA is transported into secretio...
Clonality	Monoclonal
Clone Designation	[Polyclonal]
Buffer	Delipidated, heat inactivated, lyophilized, stable whole antiserum No preservative added. Total protein and IgG concentrations in the antiserum are comparable to those of pooled normal goat serum. No foreign proteins added. Reconstitute the lyophilized a

Source	Purified free secretory component isolated from pooled rhesus monkey milk is used for immunization. Freund's complete adjuvant is used in the first step of the immunization procedure.
Application Notes	Precipitation assays. In immunoelectrophoresis use 2 μ l serum or equivalent against 120 μ l antiserum. In double radial immunodiffusion use a rosette arrangement with 10 μ l antiserum in 3 mm diameter centre well and 2 μ l serum samples (neat and serially d