

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

## Product Datasheet

### Goat anti Rabbit C3c, Clone: [Polyclonal], Monoclonal NMB-GAR/C3C

|                            |  |
|----------------------------|--|
| Article Name               | Goat anti Rabbit C3c, Clone: [Polyclonal], Monoclonal  |
| Biozol Catalog Number      | NMB-GAR/C3C  |
| Supplier Catalog Number    | GAR/C3c  |
| Alternative Catalog Number | NMB-GAR/C3C  |
| Manufacturer               | NordicMubio  |
| Host                       | Goat   |
| Category                   | Antikörper   |
| Product Description        | In immunoelectrophoresis against fresh rabbit serum, a single precipitin line is obtained in the beta-1 region representing native C3. Against serum containing partly activated C3, a precipitin line is obtained which extends from the beta-1 into the ...  |
| Clonality                  | Monoclonal   |
| Clone Designation          | [Polyclonal]   |
| UniProt                    | <a href="#">P12247</a>   |
| Buffer                     | Delipidated, heat inactivated, lyophilized, stable whole serum. No preservative added. Total protein and IgG concentrations in the antiserum are comparable to those of pooled normal goat serum. No foreign proteins added. Reconstitute the lyophilized anti |
| Source                     | C3 is the most abundant complement protein in rabbit serum. Its biological function strongly resembles that of C3 in man and other laboratory animal species. It has a central role in the activation system being common to both pathways. Activation of C3 i |

|                   |  |
|-------------------|--|
| Formula           | Delipidated, heat inactivated, lyophilized, stable whole serum. No preservative added. Total protein and IgG concentrations in the antiserum are comparable to those of pooled normal goat serum. No foreign proteins added.   |
| Application Notes | Precipitation assays. In immunoelectrophoresis use 2 $\mu$ l rabbit plasma or equivalent against 120 $\mu$ l antiserum. In double radial immunodiffusion use a rosette arrangement with 10 $\mu$ l antiserum in 3 mm diameter center well and 2 $\mu$ l plasma samples (neat and s |