

Diagnostica Vertrieb GmbH, Leipziger Straße 4

85386 Eching, Germany

Telephone: +49 (0)89 3799666-6 | **Fax:** +49 (0)89 3799666-99

E-Mail: info@biozol.de

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Goat anti Rat IgE (Fc specific), Clone: [Polyclonal], Monoclonal NMB-GARA/IGE(FC)

Article Name	Goat anti Rat IgE (Fc specific), Clone: [Polyclonal], Monoclonal
Biozol Catalog Number	NMB-GARA/IGE(FC)
Supplier Catalog Number	GARa/IgE(Fc)
Alternative Catalog Number	NMB-GARA/IGE(FC)
Manufacturer	NordicMubio
Host	Goat
Category	Antikörper
Product Description	The reactivity of the antiserum is restricted to the Fc part of the IgE molecule, which represents strict isotype (class) specificity. In immunoelectrophoresis and radial immunodiffusion, using various antiserum concentrations against appropriate con
Clonality	Monoclonal
Clone Designation	[Polyclonal]
Buffer	Delipidated, heat inactivated, lyophilized, stable whole antiserum No preservative added Total protein and IgG concentrations in the antiserum are comparable to those of pooled normal goat serum. No foreign proteins added. Reconstitute the lyophilized an
Source	Highly purified homogenous IgE isolated from rat serum. Freunds complete adjuvant is used in the first step of the immunization procedure.

Formula	Delipidated, heat inactivated, lyophilized, stable whole antiserum No preservative added Total protein and IgG concentrations in the antiserum are comparable to those of pooled normal goat serum. No foreign proteins added.
Application Notes	Precipitation assays. In immunoelectrophoresis use 2 μ l serum or equivalent against 120 μ l antiserum. In double radial immunodiffusion (Ouchterlony) use a rosette arrangement with 10 μ l antiserum in 3 mm diameter center well and 2 μ l serum samples (neat