

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

## Product Datasheet

### **Goat anti Rat IgE (Fc specific), conjugated with TRITC, Clone: [Polyclonal], Monoclonal NMB-GARA/IGE(FC)/TRITC**

|                            |  |
|----------------------------|--|
| Article Name               | Goat anti Rat IgE (Fc specific), conjugated with TRITC, Clone: [Polyclonal], Monoclonal  |
| Biozol Catalog Number      | NMB-GARA/IGE(FC)/TRITC   |
| Supplier Catalog Number    | GARa/IgE(Fc)/TRITC   |
| Alternative Catalog Number | NMB-GARA/IGE(FC)/TRITC   |
| Manufacturer               | NordicMubio  |
| Host                       | Goat   |
| Category                   | Antikörper   |
| Conjugation                | TRITC  |
| Product Description        | The reactivity of the antiserum is directed to the Fc subunits of the IgE molecule as tested by gelimmunodiffusion techniques against rat sera and a panel of purified homogenous immunoglobulins. A reaction of full identity is obtained with serum cont...  |
| Clonality                  | Monoclonal   |
| Clone Designation          | [Polyclonal]   |
| Buffer                     | TRITC-coupled purified hyperimmune goat IgG lyophilized from a solution in phosphate buffered saline (PBS, pH 7.2). No preservative added It is reconstituted by adding 1 ml sterile distilled water, spun down to remove insoluble particles, divided into sm |

|                   |   |
|-------------------|---|
| Source            | Purified homogenous IgE isolated from pooled rat serum. Freund's complete adjuvant is used in the first step of the immunization procedure. |
| Formula           | TRITC-coupled purified hyperimmune goat IgG lyophilized from a solution in phosphate buffered saline (PBS, pH 7.2). No preservative added   |
| Application Notes | ELISA, Immunocytochemistry, Immunohistochemistry (frozen), (In)direct immunofluorescence.   |