

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Goat anti Rat IgG1 IgG2a IgG2b IgG2c IgA IgM (heavy and light chains), conjugated with FITC, Clone: [Polyclonal], Monoclonal NMB-GARA/IG/FITC

| Article Name | Goat anti Rat IgG1 IgG2a IgG2b IgG2c IgA IgM (heavy and light chains), conjugated with FITC, Clone: [Polyclonal], Monoclonal |
|----------------------------|---|
| Biozol Catalog Number | NMB-GARA/IG/FITC |
| Supplier Catalog Number | GARa/Ig/FITC |
| Alternative Catalog Number | NMB-GARA/IG/FITC |
| Manufacturer | NordicMubio |
| Host | Goat |
| Category | Antikörper |
| Conjugation | FITC |
| Product Description | The reactivity of the antiserum is directed to the major isotypes of the rat immunoglobulin system (classes and both light chain types) including antibodies to common determinants, to class and to the surface determinants of the common Fab portion, a |
| Clonality | Monoclonal |
| Clone Designation | [Polyclonal] |
| Buffer | FITC-coupled purified hyperimmune goat IgG lyophilized from a solution in phosphate buffered saline (PBS, pH 7.2). No preservative added, as it may interfere with the antibody activity. It is reconstituted by adding 1 ml sterile distilled water. |

| Source | Purified polyclonal rat IgG, and homogenous IgA and IgM isolated from rat serum. Freunds complete adjuvant is used in the first step of the immunization procedure. |
|-------------------|---|
| Formula | FITC-coupled purified hyperimmune goat IgG lyophilized from a solution in phosphate buffered saline (PBS, pH 7.2). No preservative added, as it may interfere with the antibody activity. |
| Application Notes | ELISA,Immunocytochemistry,Immunohistochemistry (frozen),(In)direct immunofluorescence. |