

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Goat anti Rat IgG2c (subclass specific), conjugated with Biotin, Clone: [Polyclonal], Monoclonal NMB-GARA/IGG2C/BIO

| | |
|----------------------------|--|
| Article Name | Goat anti Rat IgG2c (subclass specific), conjugated with Biotin, Clone: [Polyclonal], Monoclonal |
| Biozol Catalog Number | NMB-GARA/IGG2C/BIO |
| Supplier Catalog Number | GARa/IgG2c/Bio |
| Alternative Catalog Number | NMB-GARA/IGG2C/BIO |
| Manufacturer | NordicMubio |
| Host | Goat |
| Category | Antikörper |
| Conjugation | Biotin |
| Product Description | The reactivity of the antiserum is directed to the subclass IgG2c. It does not react with other subclasses of IgG, IgG/Fab fragments, IgM and IgA or any non-Ig protein in rat serum, as tested by immunoelectrophoresis and double radial immunodiffusion... |
| Clonality | Monoclonal |
| Clone Designation | [Polyclonal] |
| Buffer | Biotin-coupled purified hyperimmune goat IgG lyophilized from a solution in phosphate buffered saline (PBS, pH 7.2). No preservative added, as it may interfere with the antibody activity. It is reconstituted by adding 1 ml sterile distilled water, spun d |

| | |
|-------------------|---|
| Source | Pools of purified homogenous IgG2c isolated from rat serum. Freund's complete adjuvant is used in the first step of the immunization procedure. |
| Formula | Biotin-coupled purified hyperimmune goat IgG lyophilized from a solution in phosphate buffered saline (PBS, pH 7.2). No preservative added, as it may interfere with the antibody activity. |
| Application Notes | ELISA, Immunocytochemistry, Immunohistochemistry (paraffin), Dot blot, Immunoblotting. |