

Diagnostica Vertrieb GmbH, Leipziger Straße 4

85386 Eching, Germany

**Telephone:** +49 (0)89 3799666-6 | **Fax:** +49 (0)89 3799666-99

E-Mail: info@biozol.de

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

## **Product Datasheet**

## Goat anti Rat IgM (Fc specific), conjugated with TRITC, Clone: [Polyclonal], Monoclonal NMB-GARA/IGM(FC)/TRITC

Article Name	Goat anti Rat IgM (Fc specific), conjugated with TRITC, Clone: [Polyclonal], Monoclonal
Biozol Catalog Number	NMB-GARA/IGM(FC)/TRITC
Supplier Catalog Number	GARa/IgM(Fc)/TRITC
Alternative Catalog Number	NMB-GARA/IGM(FC)/TRITC
Manufacturer	NordicMubio
Host	Goat
Category	Antikörper
Conjugation	TRITC
Product Description	The reactivity of the antiserum is directed to the Fc subunit of the IgM molecule which expresses strict isotypic (class) specificity. It does not react with purified IgG, IgG/Fab fragments and IgA or any non-Ig protein in rat serum, as tested by imm
Clonality	Monoclonal
Clone Designation	[Polyclonal]
Buffer	TRITC-coupled purified hyperimmune goat IgG lyophilized from a solution in phosphate buffered saline (PBS, pH 7.2). No preservative added, as it may interfere with the antibody activity. It is reconstituted by adding 1 ml sterile distilled water, spun do

Source	Purified pools of homogenous IgM isolated from rat serum. Freunds complete adjuvant is used in the first step of the immunization procedure.
Formula	TRITC-coupled purified hyperimmune goat IgG lyophilized from a solution in phosphate buffered saline (PBS, pH 7.2). No preservative added, as it may interfere with the antibody activity.
Application Notes	ELISA,Immunocytochemistry,Immunohistochemistry (frozen),(In)direct immunofluorescence.