

Diagnostica Vertrieb GmbH, Leipziger Straße 4

85386 Eching, Germany

Telephone: +49 (0)89 3799666-6 | **Fax:** +49 (0)89 3799666-99

E-Mail: info@biozol.de

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Goat anti Swine haptoglobin, Clone: [Polyclonal], Monoclonal NMB-GASW/HP

| Article Name | Goat anti Swine haptoglobin, Clone: [Polyclonal], Monoclonal |
|----------------------------|--|
| Biozol Catalog Number | NMB-GASW/HP |
| Supplier Catalog Number | GASw/Hp |
| Alternative Catalog Number | NMB-GASW/HP |
| Manufacturer | NordicMubio |
| Host | Goat |
| Category | Antikörper |
| Application | IP |
| Species Reactivity | Porcine |
| Product Description | The reactivity of the antiserum is restricted to haptoglobin. In immunoelectrophoresis and radial immunodiffusion (Ouchterlony), using various antiserum concentrations against fresh normal swine plasma a single precipitin line is obtained which shows |
| Clonality | Monoclonal |
| Clone Designation | [Polyclonal] |
| UniProt | Q8SPS7 |
| Buffer | Delipidated, heat inactivated, lyophilized, stable whole antiserum No preservative added Total protein and IgG concentrations in the antiserum are comparable to those of pooled normal goat serum. No foreign proteins added. Reconstitute the lyophilized an |

| Source | Haptoglobin is an acute phase protein synthesized in the liver. In immunoelectrophoresis it has an alpha-2 mobility. It forms a group of heterogeneous proteins consisting of four peptide chains, two alpha and two beta chains. The molecular weight is abou |
|-------------------|--|
| Application Notes | Precipitation assays. In immunoelectrophoresis use 2 ml or equivalent against 120 ml antiserum. In double radial immunodiffusion (Ouchterlony) use a rosette arrangement with 10 ml in a 3 mm diameter center well and 2 ml serum samples (neat and serially d |