

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

## Product Datasheet

### Goat anti Swine IgA (Fc specific), Clone: [Polyclonal], Monoclonal NMB-GASW/IGA(FC)

|                            |  |
|----------------------------|--|
| Article Name               | Goat anti Swine IgA (Fc specific), Clone: [Polyclonal], Monoclonal   |
| Biozol Catalog Number      | NMB-GASW/IGA(FC)   |
| Supplier Catalog Number    | GASw/IgA(Fc)   |
| Alternative Catalog Number | NMB-GASW/IGA(FC)   |
| Manufacturer               | NordicMubio  |
| Host                       | Goat   |
| Category                   | Antikörper   |
| Product Description        | The reactivity of the antiserum is restricted to the Fc part of the IgA molecule. In immunoelectrophoresis and radial immunodiffusion, using various antiserum concentrations against normal swine serum a single precipitin line is obtained which shows ...  |
| Clonality                  | Monoclonal   |
| Clone Designation          | [Polyclonal]   |
| Buffer                     | Delipidated, heat inactivated, lyophilized, stable whole antiserum No preservative added. Total protein and IgG concentrations in the antiserum are comparable to those of pooled normal goat serum. No foreign proteins added. Reconstitute the lyophilized a |
| Source                     | Highly purified normal IgA isolated from pooled swine serum. Freund's complete adjuvant is used in the first step of the immunization procedure.   |

|                   |   |
|-------------------|---|
| Formula           | Delipidated, heat inactivated, lyophilized, stable whole antiserum No preservative added. Total protein and IgG concentrations in the antiserum are comparable to those of pooled normal goat serum. No foreign proteins added.                               |
| Application Notes | Precipitation assays. In immunoelectrophoresis use 2 µl serum or equivalent against 120 µl antiserum. In double radial immunodiffusion (Ouchterlony) use a rosette arrangement with 10 µl antiserum in 3 mm diameter center well and 2 µl serum samples (neat |