

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Goat anti Swine IgG (heavy and light chains), conjugated with Horseradish peroxidase, Clone: [Polyclonal], HRP, Monoclonal NMB-GASW/IGG(H+L)/PO

| | |
|----------------------------|--|
| Article Name | Goat anti Swine IgG (heavy and light chains), conjugated with Horseradish peroxidase, Clone: [Polyclonal], HRP, Monoclonal |
| Biozol Catalog Number | NMB-GASW/IGG(H+L)/PO |
| Supplier Catalog Number | GASw/IgG(H+L)/PO |
| Alternative Catalog Number | NMB-GASW/IGG(H+L)/PO |
| Manufacturer | NordicMubio |
| Host | Goat |
| Category | Antikörper |
| Conjugation | HRP |
| Product Description | The reactivity of the antiserum is directed to the Fc and Fab subunits of the IgG molecule. It includes a certain degree of reactivity with other immunoglobulins via the common Fab portion. It does not react with any non-Ig protein in swine serum, as... |
| Clonality | Monoclonal |
| Clone Designation | [Polyclonal] |
| Buffer | Peroxidase-coupled purified hyperimmune goat IgG lyophilized from a solution in phosphate buffered saline (PBS, pH 7.2). No preservative added, as it may interfere with the antibody activity. It is reconstituted by adding 1 ml sterile distilled water, sp |

| | |
|-------------------|---|
| Source | Purified normal IgG isolated from pooled swine serum. Freund's complete adjuvant is used in the first step of the immunization procedure. |
| Formula | Peroxidase-coupled purified hyperimmune goat IgG lyophilized from a solution in phosphate buffered saline (PBS, pH 7.2). No preservative added, as it may interfere with the antibody activity. |
| Application Notes | ELISA, Immunocytochemistry, Immunohistochemistry (paraffin), Dot blot, Immunoblotting. |