

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Goat anti Turkey IgG (heavy and light chains), conjugated with Biotin, Clone: [Polyclonal], Monoclonal NMB-GATU/IGG(H+L)/BIO

| | |
|----------------------------|--|
| Article Name | Goat anti Turkey IgG (heavy and light chains), conjugated with Biotin, Clone: [Polyclonal], Monoclonal |
| Biozol Catalog Number | NMB-GATU/IGG(H+L)/BIO |
| Supplier Catalog Number | GATu/IgG(H+L)/Bio |
| Alternative Catalog Number | NMB-GATU/IGG(H+L)/BIO |
| Manufacturer | NordicMubio |
| Host | Goat |
| Category | Antikörper |
| Conjugation | Biotin |
| Product Description | The reactivity of the antiserum is directed to the Fc and Fab subunits of the IgG molecule. It includes a certain degree of reactivity with other immunoglobulins via the common Fab portion. It does not react with any non-Ig protein in turkey serum, a... |
| Clonality | Monoclonal |
| Clone Designation | [Polyclonal] |
| Buffer | Biotin-coupled purified hyperimmune goat IgG lyophilized from a solution in phosphate buffered saline. No preservative added. It is reconstituted by adding 1 ml sterile distilled water, spun down to remove insoluble particles, divided into small aliquots |

| | |
|-------------------|--|
| Source | Purified normal IgG isolated from pooled turkey serum. Freund's complete adjuvant is used in the first step of the immunization procedure. |
| Formula | Biotin-coupled purified hyperimmune goat IgG lyophilized from a solution in phosphate buffered saline. No preservative added. |
| Application Notes | ELISA, Immunocytochemistry, Immunohistochemistry (paraffin), Dot blot, Immunoblotting. |