

Diagnostica Vertrieb GmbH, Leipziger Straße 4

85386 Eching, Germany

Telephone: +49 (0)89 3799666-6 | **Fax:** +49 (0)89 3799666-99

E-Mail: info@biozol.de

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Rabbit anti Goat IgG (Fc specific), Clone: [Polyclonal], Monoclonal NMB-RAG/IGG(FC)

Article Name	Rabbit anti Goat IgG (Fc specific), Clone: [Polyclonal], Monoclonal
Biozol Catalog Number	NMB-RAG/IGG(FC)
Supplier Catalog Number	RAG/IgG(Fc)
Alternative Catalog Number	NMB-RAG/IGG(FC)
Manufacturer	NordicMubio
Host	Rabbit
Category	Antikörper
Product Description	The reactivity of the antiserum is restricted to the Fc part of the IgG molecule. In immunoelectrophoresis and radial immunodiffusion, using various antiserum concentrations against normal goat serum a single precipitin line is obtained which shows a
Clonality	Monoclonal
Clone Designation	[Polyclonal]
Buffer	Delipidated, heat inactivated, lyophilized, stable whole antiserum. No preservative added. Total protein and IgG concentrations in the antiserum are comparable to those of pooled normal rabbit serum. No foreign proteins added. Reconstitute the lyophilize
Source	Highly purified normal IgG isolated from pooled goat serum. Freunds complete adjuvant is used in the first step of the immunization procedure.

Formula	Delipidated, heat inactivated, lyophilized, stable whole antiserum. No preservative added. Total protein and IgG concentrations in the antiserum are comparable to those of pooled normal rabbit serum. No foreign proteins added.
Application Notes	Precipitation assays. In immunoelectrophoresis use 2 μ l serum or equivalent against 120 μ l antiserum. In double radial immunodiffusion (Ouchterlony) use a rosette arrangement with 10 μ l antiserum in 3 mm diameter center well and 2 μ l serum samples (neat