

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Rabbit anti Goat IgG (heavy and light chains), Clone: [Polyclonal], Monoclonal NMB-RAG/IGG(H+L)

Article Name	Rabbit anti Goat IgG (heavy and light chains), Clone: [Polyclonal], Monoclonal
Biozol Catalog Number	NMB-RAG/IGG(H+L)
Supplier Catalog Number	RAG/IgG(H+L)
Alternative Catalog Number	NMB-RAG/IGG(H+L)
Manufacturer	NordicMubio
Host	Rabbit
Category	Antikörper
Product Description	The reactivity of the antiserum is directed to the Fc and Fab subunits of the IgG molecule. It includes a certain degree of reactivity with other immunoglobulins via the common Fab portion. It does not react with any non-Ig protein in goat serum, as ...
Clonality	Monoclonal
Clone Designation	[Polyclonal]
Buffer	Delipidated, heat inactivated lyophilized stable whole serum. No preservative added, as it may interfere with the antibody activity. No foreign protein added. Total protein and IgG concentration in the antiserum are comparable to those of pooled rabbit s
Source	Purified normal IgG isolated from pooled goat serum. Freund's complete adjuvant is used in the first step of the immunization procedure.

Formula	Delipidated, heat inactivated lyophilized stable whole serum. No preservative added, as it may interfere with the antibody activity. No foreign protein added. Total protein and IgG concentration in the antiserum are comparable to those of pooled rabbit s
Application Notes	Precipitation assays. In immunoelectrophoresis use 2 µl or equivalent against 120 µl antiserum. In double radial immunodiffusion (Ouchterlony) use a rosette arrangement with 10 µl antiserum in a 3 mm diameter centre well and 2 µl serum samples (neat and