

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

## Product Datasheet

### **Rabbit anti Goat Ig light chains kappa and lambda (free and bound), Clone: [Polyclonal], Monoclonal NMB-RAG/L(SD+HD)**

Article Name	Rabbit anti Goat Ig light chains kappa and lambda (free and bound), Clone: [Polyclonal], Monoclonal
Biozol Catalog Number	NMB-RAG/L(SD+HD)
Supplier Catalog Number	RAG/L(SD+HD)
Alternative Catalog Number	NMB-RAG/L(SD+HD)
Manufacturer	NordicMubio
Host	Rabbit
Category	Antikörper
Product Description	As tested in immunoelectrophoresis and double radial immunodiffusion against pooled normal goat serum and purified IgG fractions and light chain preparations, the antiserum reacts with determinants exposed on the surface (SD) and the Fab portion of a...
Clonality	Monoclonal
Clone Designation	[Polyclonal]
Buffer	Delipidated, heat inactivated lyophilized stable whole serum. No preservative added, as it may interfere with the antibody activity. No foreign protein added. Total protein and IgG concentration in the antiserum are comparable to those of pooled rabbit s
Source	Purified light chains isolated from polyclonal goat IgG. Freund's complete adjuvant is used in the first step of the immunization procedure.

Formula	Delipidated, heat inactivated lyophilized stable whole serum. No preservative added, as it may interfere with the antibody activity. No foreign protein added. Total protein and IgG concentration in the antiserum are comparable to those of pooled rabbit s
Application Notes	Precipitation assays. In immunoelectrophoresis use 2 µl or equivalent against 120 µl antiserum. In double radial immunodiffusion (Ouchterlony) use a rosette arrangement with 10 µl antiserum in a 3 mm diameter centre well and 2 µl serum samples (neat and