

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Rabbit anti Goat IgM (heavy and light chains), conjugated with FITC, Clone: [Polyclonal], Monoclonal NMB-RAG/IGM(H+L)/FITC

| | |
|----------------------------|---|
| Article Name | Rabbit anti Goat IgM (heavy and light chains), conjugated with FITC, Clone: [Polyclonal], Monoclonal |
| Biozol Catalog Number | NMB-RAG/IGM(H+L)/FITC |
| Supplier Catalog Number | RAG/IgM(H+L)/FITC |
| Alternative Catalog Number | NMB-RAG/IGM(H+L)/FITC |
| Manufacturer | NordicMubio |
| Host | Rabbit |
| Category | Antikörper |
| Conjugation | FITC |
| Product Description | The reactivity of the antiserum is directed to the Fc and Fab subunits of the IgM molecule. It includes a certain degree of reactivity with other immunoglobulins via the common Fab portion. It does not react with any non-Ig protein in sheep serum, as... |
| Clonality | Monoclonal |
| Clone Designation | [Polyclonal] |
| Buffer | FITC-coupled purified hyperimmune rabbit IgG lyophilized from a solution in phosphate buffered saline (PBS, pH 7.2). No preservative added. It is reconstituted by adding 1 ml sterile distilled water, spun down to remove insoluble particles, divided into |

| | |
|-------------------|---|
| Source | Purified normal IgM isolated from pooled goat serum. Freund's complete adjuvant is used in the first step of the immunization procedure. |
| Formula | FITC-coupled purified hyperimmune rabbit IgG lyophilized from a solution in phosphate buffered saline (PBS, pH 7.2). No preservative added. |
| Application Notes | ELISA, Immunocytochemistry, Immunohistochemistry (frozen), (In)direct immunofluorescence. |