

Diagnostica Vertrieb GmbH, Leipziger Straße 4

85386 Eching, Germany

Telephone: +49 (0)89 3799666-6 | **Fax:** +49 (0)89 3799666-99

E-Mail: info@biozol.de

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Rabbit anti Guinea Pig IgG1 (subclass specific), Clone: [Polyclonal], Monoclonal NMB-RAGP/IGG1

| Article Name | Rabbit anti Guinea Pig IgG1 (subclass specific), Clone: [Polyclonal], Monoclonal |
|----------------------------|--|
| Biozol Catalog Number | NMB-RAGP/IGG1 |
| Supplier Catalog Number | RAGp/lgG1 |
| Alternative Catalog Number | NMB-RAGP/IGG1 |
| Manufacturer | NordicMubio |
| Host | Rabbit |
| Category | Antikörper |
| Application | IP |
| Species Reactivity | Guinea pig |
| Product Description | The reactivity of the antiserum is directed to the Fc subunit of subclass IgG1. It does not react with other IgG2, IgG/Fab fragments, IgM and IgA or any non-Ig protein in guinea pig serum, as tested by immunoelectrophoresis and double radial immunodi |
| Clonality | Monoclonal |
| Clone Designation | [Polyclonal] |
| Buffer | Delipidated, heat inactivated, lyophilized whole antiserum. No preservative added, as it may interfere with the antibody activity. Total protein and IgG concentration in the antiserum are comparable to those of pooled normal rabbit serum. No foreign prot |

| Source | Pools of purified normal IgG1 isolated from pooled guinea pig serum. Freunds complete adjuvant is used in the first step of the immunization procedure. |
|-------------------|---|
| Application Notes | Precipitation assays. In immunoelectrophoresis use 2 μ l or equivalent against 120 μ l antiserum. In double radial immunodiffusion (Ouchterlony) use a rosette arrangement with 10 μ l antiserum in a 3 mm diameter centre well and 2 μ l serum samples (neat and |