

Diagnostica Vertrieb GmbH, Leipziger Straße 4

85386 Eching, Germany

**Telephone:** +49 (0)89 3799666-6 | **Fax:** +49 (0)89 3799666-99

E-Mail: info@biozol.de

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

## **Product Datasheet**

## Rabbit anti Horse IgG (heavy and light chains), Clone: [Polyclonal], Monoclonal NMB-RAHO/IGG(H+L)

| Article Name               | Rabbit anti Horse IgG (heavy and light chains), Clone: [Polyclonal], Monoclonal  |
|----------------------------|--|
| Biozol Catalog Number      | NMB-RAHO/IGG(H+L)  |
| Supplier Catalog Number    | RAHo/lgG(H+L)  |
| Alternative Catalog Number | NMB-RAHO/IGG(H+L)  |
| Manufacturer               | NordicMubio  |
| Host                       | Rabbit   |
| Category                   | Antikörper   |
| Product Description        | The reactivity of the antiserum is directed to the Fc and Fab subunits of both subclasses IgG1 and IgG2. Reactivity with other subclasses of horse IgG has not been tested. The antiserum reacts with both Fc and Fab portions of polyclonal IgG. It inclu     |
| Clonality                  | Monoclonal   |
| Clone Designation          | [Polyclonal]   |
| Buffer                     | Delipidated, heat inactivated lyophilized stable whole serum. No preservative added, as it may interfere with the antibody activity. No foreign protein added. Total protein and IgG concentration in the antiserum are comparable to those of pooled rabbit s |
| Source                     | Purified normal IgG isolated from pooled horse serum. Freunds complete adjuvant is used in the first step of the immunization procedure.   |

| Formula           | Delipidated, heat inactivated lyophilized stable whole serum. No preservative added, as it may interfere with the antibody activity. No foreign protein added. Total protein and IgG concentration in the antiserum are comparable to those of pooled rabbit s                    |
|-------------------|---|
| Application Notes | Precipitation assays. In immunoelectrophoresis use 2 $\mu$ l or equivalent against 120 $\mu$ l antiserum. In double radial immunodiffusion (Ouchterlony) use a rosette arrangement with 10 $\mu$ l antiserum in a 3 mm diameter centre well and 2 $\mu$ l serum samples (neat and |