

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

## Product Datasheet

### Rabbit anti Human IgG Fab, Clone: [Polyclonal], Monoclonal NMB-RAHU/FAB

|                            |  |
|----------------------------|--|
| Article Name               | Rabbit anti Human IgG Fab, Clone: [Polyclonal], Monoclonal   |
| Biozol Catalog Number      | NMB-RAHU/FAB   |
| Supplier Catalog Number    | RAHu/Fab   |
| Alternative Catalog Number | NMB-RAHU/FAB   |
| Manufacturer               | NordicMubio  |
| Host                       | Rabbit   |
| Category                   | Antikörper   |
| Product Description        | The antiserum is reacting with the Fab subunit of intact IgG, IgA and IgM and other Ig classes of both light chain types, with their Fab or F(ab)2 subunits and with free light chains of kappa and lambda type as tested in immunoelectrophoresis and dou...  |
| Clonality                  | Monoclonal   |
| Clone Designation          | [Polyclonal]   |
| Buffer                     | Delipidated, heat inactivated, lyophilized, stable whole serum. No preservative added. Total protein and IgG concentrations in the antiserum are comparable to those of normal pooled rabbit serum. No foreign proteins added. It is reconstituted by adding 1 |
| Source                     | Purified Fab of normal human IgG. Freund's complete adjuvant is used in the first step of the immunization procedure.  |
| Formula                    | Delipidated, heat inactivated, lyophilized, stable whole serum. No preservative added. Total protein and IgG concentrations in the antiserum are comparable to those of normal pooled rabbit serum. No foreign proteins added.                                 |

Application Notes

Precipitation assays. In immunoelectrophoresis use 2  $\mu$ l or equivalent against 120  $\mu$ l antiserum. In double radial immunodiffusion (Ouchterlony) use a rosette arrangement with 10  $\mu$ l antiserum in a 3 mm diameter centre well and 2  $\mu$ l serum samples (neat and