

Diagnostica Vertrieb GmbH, Leipziger Straße 4

85386 Eching, Germany

**Telephone:** +49 (0)89 3799666-6 | **Fax:** +49 (0)89 3799666-99

E-Mail: info@biozol.de

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

## **Product Datasheet**

## Rabbit anti Human IgG IgA IgM IgD IgE (heavy and light chains), Clone: [Polyclonal], Monoclonal NMB-RAHU/IG(GAMDE)

| Article Name               | Rabbit anti Human IgG IgA IgM IgD IgE (heavy and light chains),<br>Clone: [Polyclonal], Monoclonal  |
|----------------------------|---|
| Biozol Catalog Number      | NMB-RAHU/IG(GAMDE)  |
| Supplier Catalog Number    | RAHu/Ig(GAMDE)  |
| Alternative Catalog Number | NMB-RAHU/IG(GAMDE)  |
| Manufacturer               | NordicMubio   |
| Host                       | Rabbit  |
| Category                   | Antikörper  |
| Product Description        | The reactivity of the antiserum is directed to the heavy and light chains of the immunoglobulin isotypes. It does not react with any non-Ig protein in human serum, as tested by immunoelectrophoresis and double radial immunodiffusion. In immunoelectro    |
| Clonality                  | Monoclonal  |
| Clone Designation          | [Polyclonal]  |
| Buffer                     | Delipidated, heat inactivated, lyophilized stable whole serum. No preservative added, as it may interfere with the antibody activity. No foreign protein added. Total protein and IgG concentration in the antiserum are comparable to those of pooled rabbit |
| Source                     | Purified normal Ig fractions isolated from pooled human serum.<br>Freunds complete adjuvant is used in the first step of the<br>immunization procedure.   |

| Formula           | Delipidated, heat inactivated, lyophilized stable whole serum. No preservative added, as it may interfere with the antibody activity. No foreign protein added. Total protein and IgG concentration in the antiserum are comparable to those of pooled rabbit                     |
|-------------------|---|
| Application Notes | Precipitation assays. In immunoelectrophoresis use 2 $\mu$ l or equivalent against 120 $\mu$ l antiserum. In double radial immunodiffusion use a rosette arrangement with 10 $\mu$ l antiserum in a 3 mm diameter centre well and 2 $\mu$ l serum samples (neat and diluted) in 2 |