

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Rabbit anti Human lactoferrin, Clone: [Polyclonal], Monoclonal NMB-RAHU/LFR

| | |
|----------------------------|--|
| Article Name | Rabbit anti Human lactoferrin, Clone: [Polyclonal], Monoclonal |
| Biozol Catalog Number | NMB-RAHU/LFR |
| Supplier Catalog Number | RAHu/Lfr |
| Alternative Catalog Number | NMB-RAHU/LFR |
| Manufacturer | NordicMubio |
| Host | Rabbit |
| Category | Antikörper |
| Product Description | In immunoelectrophoresis against human milk a single precipitin line is obtained. The antiserum does not react with any other protein component of human serum or plasma. In precipitating techniques as immunoelectrophoresis, single and double radial i... |
| Clonality | Monoclonal |
| Clone Designation | [Polyclonal] |
| UniProt | P02788 |
| Buffer | Delipidated, heat inactivated lyophilized whole serum. No preservative added. Total protein and IgG concentration in the antiserum are comparable to those of pooled normal rabbit serum. No foreign proteins added. Reconstitute the lyophilized antiserum by |
| Source | Exocrine organs produce various secretions, each with its characteristic function. Proteins found in secretions may be divided into two groups: those specific for the particular secretion, and plasma proteins independent of the type of exocrine cells. La |

| | |
|-------------------|---|
| Formula | Delipidated, heat inactivated lyophilized whole serum. No preservative added. Total protein and IgG concentration in the antiserum are comparable to those of pooled normal rabbit serum. No foreign proteins added. |
| Application Notes | Precipitation assays. In immunoelectrophoresis use 2 μ l human serum against 120 μ l antiserum. In double radial immunodiffusion (Ouchterlony) use a rosette arrangement with 10 μ l antiserum in a 3 mm diameter center well and 10 μ l of the antiserum in 3 mm |