

Diagnostica Vertrieb GmbH, Leipziger Straße 4

85386 Eching, Germany

Telephone: +49 (0)89 3799666-6 | **Fax:** +49 (0)89 3799666-99

E-Mail: info@biozol.de

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Rabbit anti Human Iysozyme, conjugated with Horseradish peroxidase, Clone: [Polyclonal], HRP, Monoclonal NMB-RAHU/LYS/PO

Article Name	Rabbit anti Human lysozyme, conjugated with Horseradish peroxidase, Clone: [Polyclonal], HRP, Monoclonal
Biozol Catalog Number	NMB-RAHU/LYS/PO
Supplier Catalog Number	RAHu/Lys/PO
Alternative Catalog Number	NMB-RAHU/LYS/PO
Manufacturer	NordicMubio
Host	Rabbit
Category	Antikörper
Species Reactivity	Human
Immunogen	Highly purified recombinant human lysozyme was used as immunogen.
Conjugation	HRP
Product Description	The reactivity of the antiserum is restricted to lysozyme. In immunoelectrophoresis and radial immunodiffusion (Ouchterlony), using various antiserum concentrations against normal human milk and tears a single precipitin line is obtained which shows
Clonality	Monoclonal
Clone Designation	[Polyclonal]
UniProt	P61626

Buffer	Horseradish peroxidase-coupled purified hyperimmune IgG lyophilized from a solution in phosphate buffered saline (PBS, pH 7.2) No preservative added, as it may interfere with the antibody activity.
Source	Lysozyme, a bacteriolytic enzyme, has a molecular weight of about 15,000. Its concentration in plasma is low, in milk the concentration is reported to be between 5 and 220 mg/ml. Also tears contain a considerable amount of lysozyme. Highly purified recom
Formula	Horseradish peroxidase-coupled purified hyperimmune IgG Iyophilized from a solution in phosphate buffered saline (PBS, pH 7.2) No preservative added, as it may interfere with the antibody activity.
Application Notes	ELISA,Immunocytochemistry,Immunohistochemistry (paraffin),Dot blot,Immunoblotting. The lyophilized conjugate is shipped at ambient temperature and may be stored at +4C, prolonged storage at or below -20C. It is reconstituted by adding 1 ml sterile di s