

Diagnostica Vertrieb GmbH, Leipziger Straße 4

85386 Eching, Germany

Telephone: +49 (0)89 3799666-6 | **Fax:** +49 (0)89 3799666-99

E-Mail: info@biozol.de

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Rabbit anti Human secretory component (free and bound), Clone: [Polyclonal], Monoclonal NMB-RAHU/SC

Article Name	Rabbit anti Human secretory component (free and bound), Clone: [Polyclonal], Monoclonal
Biozol Catalog Number	NMB-RAHU/SC
Supplier Catalog Number	RAHu/SC
Alternative Catalog Number	NMB-RAHU/SC
Manufacturer	NordicMubio
Host	Rabbit
Category	Antikörper
Application	ELISA, IP
Species Reactivity	Human
Product Description	Tested in immunoelectrophoresis, double radial immunodiffusion and ELISA against a panel of appropriate secretions and purified Ig isotypes. The antiserum reacts with both bound secretory component (secretory IgA) and with the free SC present in huma
Clonality	Monoclonal
Clone Designation	[Polyclonal]
UniProt	P01833
Buffer	Delipidated, heat inactivated, lyophilized, stable whole antiserum. No preservative added. Total protein and IgG concentrations in the antiserum are comparable to those of pooled normal rabbit serum. No foreign proteins added. Reconstitute the lyophilize

Source	Secretory component is present in human secretions bound to secretory IgA (sIgA) and in free form. Secretory IgA (sIgA) functions as a dimer or polymer and accounts for almost all specific mucosal antibody activity. A molecule of sIgA is made up of two m
Formula	Delipidated, heat inactivated, lyophilized, stable whole antiserum. No preservative added. Total protein and IgG concentrations in the antiserum are comparable to those of pooled normal rabbit serum. No foreign proteins added.
Application Notes	Precipitation assays and ELISA. In immunoelectrophoresis use 2 μ l serum or equivalent against 120 μ l antiserum. In double radial immunodiffusion use a rosette arrangement with 10 μ l antiserum in 3 mm diameter centre well and 2 μ l serum samples (neat and